

# (12) UK Patent Application (19) GB (11) 2 275 479 (13) A

(43) Date of A Publication 31.08.1994

(21) Application No 9402090.6

(22) Date of Filing 03.02.1994

(30) Priority Data

(31) 9303912 (32) 26.02.1993 (33) GB

(71) Applicant(s)

Zeneca Limited

(Incorporated in the United Kingdom)

Imperial Chemical House, 9 Millbank, LONDON,  
SW1P 3JF, United Kingdom

(72) Inventor(s)

Peter Gregory  
Ronald Wynford Kenyon  
Roberts Bowker

(74) Agent and/or Address for Service

Roger Graham Pugsley  
Imperial Chemical Industries PLC, ICI Group Patents,  
Group Patents Services Dept, PO Box 6, Shire Park,  
Bessemer Road, WELWYN GARDEN CITY,  
Hertfordshire, AL7 1HD, United Kingdom

(51) INT CL<sup>5</sup>

C09B 67/22, C09D 11/02

(52) UK CL (Edition M)

C4P PM P110  
U1S S2239

(56) Documents Cited


EP 0565286 A EP 0559309 A EP 0534634 A  
EP 0494523 A EP 0468649 A EP 0468648 A  
EP 0468647 A EP 0356080 A US 4963189 A

(58) Field of Search

UK CL (Edition M) C4P PM  
INT CL<sup>5</sup> C09B 67/22, C09D 11/00 11/02  
ONLINE DATABASES: CA;WPI;CLAIMS

(54) Ink

(57) An ink suitable for use in ink jet printing comprising a solution of a colorant in an aqueous medium, the colorant comprising at least 80% by weight of a water-soluble black dye of Formula (1)



wherein

W is COOH  
X is H, COOH, SO<sup>3</sup>H, halo, hydroxy, nitro, cyano, C<sub>1-6</sub>-alkyl, C<sub>1-6</sub>-alkoxy, or C<sub>1-6</sub>-acylamino;  
Y is H, COOH or SO<sub>3</sub>H;  
Z is H, COOH or SO<sub>3</sub>H; and  
R is H or C<sub>1-4</sub>-alkyl;

provided that there are at least two COOH groups and that the number of COOH groups is equal to or greater than the number of SO<sub>3</sub>H groups and up to 20% of one or both of two other water-soluble dyes, one of which is a defined yellow dye and the other is a defined cyan dye.

GB 2 275 479 A

**2275479****INK**

This specification describes an invention relating to an ink suitable for use in ink jet printing and, in particular, a neutral black ink with good wet-fastness.

According to the present invention there is provided an ink comprising a solution of a colorant in an aqueous medium, the colorant comprising at least 80% by weight of a water-soluble black dye of Formula (1) as hereinafter defined and up to 20% of one or both of two other water-soluble dyes, one of which is a yellow dye of the Formula (2) as hereinafter defined and the other is a cyan dye of Formula (9) as hereinafter defined.

The colorant preferably comprises at least 85%, more preferably at least 90% and preferably up to 98% by weight of the black dye and up to 15% by weight of the yellow and/or the cyan dye, more preferably up to 10% and preferably at least 2%. One specifically preferred colorant comprises from 90% to 95% of the black dye and from 10% to 5% of the cyan dye, another specifically preferred colorant comprises from 90% to 98% of the black dye and from 10% to 2% of the yellow dye and another specifically preferred colorant comprises from 90% to 98% of the black dye and from 10% to 2% of a mixture of the yellow dye and the cyan dye. The mixture of yellow and cyan dyes preferably comprises from 90:10 to 10:90, more preferably from 80:20 to 20:80 and especially from 60:40 to 40:60 by weight of yellow dye to cyan dye.

The amount of the colorant in the ink is preferably from 0.5% to 20%, more preferably from 0.5% to 15% and especially from 1% to 5%, by weight with respect to the total weight of the ink. The aqueous medium is preferably substantially water but preferably comprises up to 20%, more preferably up to 15% and especially up to 10%, by weight of one or more water-soluble organic liquids. The ink may also comprise other conventional additives, especially those used in ink jet printing inks.

The water-soluble organic liquid(s) is conveniently selected from C<sub>1-4</sub>-alkanols, such as methanol, ethanol, n-propanol, isopropanol, n-butanol, sec-butanol, tert-butanol and isobutanol; amides such as dimethylformamide and dimethylacetamide; ketones and ketone alcohols such as acetone and diacetone alcohol; ethers such as tetrahydrofuran and dioxane; alkylene glycols or thioglycols containing a C<sub>2</sub>-C<sub>6</sub> alkylene group such as ethylene glycol, propylene glycol, butylene

glycol, pentylene glycol and hexylene glycol,, poly(alkylene-glycol)s and thioglycol)s such as diethylene glycol, thiodiglycol, polyethylene glycol and polypropylene glycol; polyols such as glycerol and 1,2,6-hexanetriol; and lower alkyl glycol and polyglycol ethers, such as 2-methoxyethanol, 2-(2-methoxyethoxy)ethanol, 2-(2-ethoxyethoxy)ethanol, 2-(2-butoxyethoxy)ethanol, 2-[2-(2-methoxyethoxy)-ethoxy]ethanol, 2-[2-(2-ethoxyethoxy)ethoxy]-ethanol; or a mixture containing two or more of the aforementioned water-soluble organic liquids.

However it is preferred that the organic liquid(s) is selected from heterocyclic ketones or sulphones, such as 2-pyrrolidone, N-methyl-2-pyrrolidone, N-(2-hydroxyethyl)-2-pyrrolidone and sulpholane, especially 2-pyrrolidone and sulpholane. Other preferred organic liquids are 2-(2-butoxyethoxy)ethanol and n-pentan-1,4-diol.

We have found that the present inks, wherein the medium contains up to 20% by weight of one or more of these preferred organic liquids, especially 2-pyrrolidone, are more stable than equivalent inks wherein the medium contains one or more of the other aforementioned organic liquids.

Examples of other ingredients which may be present in the inks are a penetrant, to assist permeation of the dye into a paper substrate, a kogation-reducing agent, to prevent or reduce the build-up of residue (koga) on the resistor surface in thermal ink jet printers, and a buffer, such as sodium borate, to stabilise the pH of the ink.

The kogation-reducing agent is preferably an oxo anion, such as described in EP 425150A. The oxo-anion may be  $C_2O_4^{2-}$ ,  $SO_3^{2-}$ ,  $SO_4^{2-}$ , molybdate,  $AsO_4^{3-}$  or more preferably a phosphate ester, a diorgano-phosphate or more especially a phosphate salt which is particularly effective in reducing kogation. Examples of phosphate salts are dibasic phosphate ( $HPO_4^{2-}$ ), monobasic phosphates ( $H_2PO_4^-$ ) and polyphosphates ( $P_2O_7^{4-}$ ). The selection of the counterion is not believed to be critical and examples include alkali metals, ammonium and alkylammonium cations. The amount of kogation-reducing agent in the ink is preferably from 0.001% to 15%, based on oxo-anion, and more preferably from 0.01% to 1%, by weight with respect to the total ink.

A further aspect of the present invention provides a process for printing a substrate with an ink as hereinbefore defined by ink jet printing.

In ink jet printing an ink is formed into small droplets by ejection from a reservoir through a small orifice so that the droplets of ink are directed at a substrate. Suitable ink jet printing processes for the present ink are piezoelectric ink jet printing, and more  
 5 especially thermal ink jet printing. In thermal ink jet printing, programmed pulses of heat are applied to the ink by means of a resistor, adjacent to the orifice during relative movement between the substrate and the reservoir.


Preferred substrates include cellulosic sheet materials, such as  
 10 clear cellulosic sheets suitable for projector slides and more preferably paper, especially plain paper, which may be acidic, alkaline or neutral.

According to a further aspect of the present invention there is provided a cellulosic substrate, especially paper or a projector slide  
 15 printed with an ink according to the invention.

Although the ink may contain other dyes than those hereinbefore defined, it preferably free from such other dyes as the ink containing only the hereinbefore defined dyes in the preferred proportions has a particularly attractive neutral black shade and especially high wet-  
 20 fastness on cellulosic substrates.

#### Black Dye

The black dye, in the free acid form, has the Formula (1):



30 wherein

- W is COOH
- X is H, COOH, SO<sup>3</sup>H, halo, hydroxy, nitro, cyano, C<sub>1-6</sub>-alkyl, C<sub>1-6</sub>-alkoxy, or C<sub>1-6</sub>-acylamino;
- Y is H, COOH or SO<sub>3</sub>H;
- 35 Z is H, COOH or SO<sub>3</sub>H; and
- R is H or C<sub>1-4</sub>-alkyl;

provided that there are at least two COOH groups and that the number of COOH groups is equal to or greater than the number of SO<sub>3</sub>H groups.

The group X is preferably selected from H; COOH; SO<sub>3</sub>H; halo especially chloro or bromo; hydroxy; nitro; cyano; C<sub>1-4</sub>-alkyl especially methyl; C<sub>1-4</sub>-alkoxy especially methoxy; and C<sub>1-4</sub>-alkylcarbonylamino; especially acetamido. However X is more preferably H or COOH, and especially COOH.

Where X is H, the COOH group on Ring A represented by W may be in the ortho, meta or para position with respect to the azo group but is preferably in the meta or para position. However, it is preferred that X is COOH and that the two COOH groups are in the 3,4-, 2,5- and especially the 3,5- positions on Ring A.

The black dye preferably contains from 2 to 4 COOH groups and preferably not more than two SO<sub>3</sub>H groups, i.e. not more than one of X, Y and Z is SO<sub>3</sub>H, i.e. there are preferably the same number of COOH and SO<sub>3</sub>H groups or an excess of 1 or 2 COOH groups.

The group R is preferably H or methyl, and more especially H.

Preferred diazo components from which Ring A is derivable are, 2-aminobenzoic acid, 3-aminobenzoic acid, 4-aminobenzoic acid, 2-aminophthalic acid, 3-aminophthalic acid, 4-aminoisophthalic acid, 2-aminoisophthalic acid, 2-aminoterephthalic acid and especially 5-aminoisophthalic acid,

Preferred couplers from which the central group B is derivable are, 1,6-Cleves acid, 1,7-Cleves acid, mixed Cleves acids, 8-amino-2-naphthoic acid, 5-amino-2-naphthoic acid and 5-amino-2,3-dicarboxy-naphthalene and especially 1-naphthylamine.

Preferred couplers from which the third group C is derivable are, N-methyl-Gamma acid, N-butyl-Gamma acid and especially Gamma acid.

Suitable black dyes of Formula (1)<sup>+</sup>, in the ammonium (\*NH<sub>4</sub>) salt form, in which R is H are set out in the following table:

| Dye | X      | W      | Y                 | Z;                 | Dye | X      | W      | Y    | Z     |
|-----|--------|--------|-------------------|--------------------|-----|--------|--------|------|-------|
| B1  | 3-COOH | 5-COOH | H                 | H;                 | B8  | 3-COOH | 5-COOH | COOH | H;    |
| B2  | 3-COOH | 5-COOH | H                 | SO <sub>3</sub> H; | B9  | 3-COOH | 5-COOH | H    | COOH; |
| B3  | 3-COOH | 5-COOH | SO <sub>3</sub> H | H;                 | B10 | 3-COOH | 5-COOH | COOH | COOH; |
| B4  | H      | 4-COOH | H                 | COOH;              | B11 | H      | 3-COOH | COOH | H;    |
| B5  | H      | 3-COOH | H                 | COOH;              |     |        |        |      |       |
| B6  | 2-COOH | 5-COOH | H                 | H;                 | B12 | 2-COOH | 3-COOH | H    | H;    |
| B7  | 2-COOH | 4-COOH | H                 | SO <sub>3</sub> H; | B13 | 3-COOH | 4-COOH | H    | H;    |


Preferred black dyes are Dye B2 and, more especially, Dye B1.

The black dye is described in greater detail in EP 356,080 the contents of which are incorporated herein by reference.

# Yellow Dye

The yellow dye, in the free acid form, has the structure shown in

5 Formula (2):



10

wherein:

Ar and Ar<sup>1</sup> are each independently aryl or substituted aryl providing at least one of Ar and Ar<sup>1</sup> has at least one substituent

15

selected from COOH and COSH;


R<sup>1</sup>, R<sup>2</sup>, R<sup>3</sup>, and R<sup>4</sup> are each independently H, alkyl or substituted alkyl;

L is a divalent organic linking group;


n is 0 or 1;

J and J<sup>1</sup> are each independently of formula (3), (4) or (5):


20



(3)



(4)



(5)

25

wherein

30 each R<sup>5</sup> is independently selected from H, alkyl, substituted alkyl, alkoxy, halogen, CN, ureido and NHCOR<sup>6</sup>;

R<sup>6</sup> is H, alkyl, substituted alkyl, aryl, substituted aryl, aralkyl or substituted aralkyl;




each T is independently alkyl;

35 each W is independently selected from H, CN, CONR<sup>10</sup>R<sup>11</sup>, pyridinium and COOH;

each M is an alkylene chain having 2 to 8 carbon atoms;

B is H, alkyl or COOH; and

$R^{10}$  and  $R^{11}$  are each independently H, alkyl or substituted alkyl;  
 each X is independently carbonyl or a group of the Formula (6), (7)  
 or (8):



(6)

(7)

(8)

wherein

Z is  $OR^7$ ,  $SR^7$  or  $NR^8R^9$ ;

Y is H, Cl, CN or Z;

E is Cl or CN; and

$R^7$ ,  $R^8$  and  $R^9$  are each independently H, alkenyl, substituted alkenyl,  
 alkyl, substituted alkyl, aryl, substituted aryl, aralkyl or  
 substituted aralkyl, or

$R^8$  and  $R^9$  together with the nitrogen atom to which they are attached  
 form a 5 or 6 membered ring;

provided that

- (a) if the compound of Formula (2) has no  $-SO_3H$  groups then it  
 has at least two groups selected from  $-COOH$  and  $-COSH$ ;
- (b) the compound of Formula (2) has at least as many groups  
 selected from  $-COOH$  and  $-COSH$  as  $-SO_3H$  groups; and
- (c) that the structures of the groups  $Ar-N=N-J$  and  $Ar-N=N-J^1$  are  
 such that the compound of Formula (2) is yellow.

It is preferred that the yellow dye of Formula (2) has at least as  
 many  $-COOH$  as  $-SO_3H$  groups and, if it has no  $-SO_3H$  groups that it has at  
 least two and, more preferably at least three,  $-COOH$  groups. It is  
 especially preferred that the yellow dye has up to 2  $-SO_3H$  groups and  
 from 2 to 5  $-COOH$ . It is also preferred that the yellow dye is free  
 from cellulose reactive groups.

The groups Ar and  $Ar^1$  are preferably independently selected from  
 phenyl and substituted phenyl. Each optional substituent on Ar and  $Ar^1$   
 is preferably selected from optionally substituted alkyl, especially  
 $C_{1-4}$ -alkyl; alkoxy, especially  $C_{1-4}$ -alkoxy;  $-SO_3H$ ;  $-PO_3H_2$ ;  $-COSH$ ;  $-OH$ ;  
 $-CO_2H$ ; and halogen, especially Cl or Br. It is especially preferred

that when Ar or Ar<sup>1</sup> is substituted phenyl, each substituent is independently selected from CO<sub>2</sub>H, COSH and SO<sub>3</sub>H, and more especially each substituent is CO<sub>2</sub>H. In especially preferred structures, at least one of Ar and Ar<sup>1</sup> has at least one -COOH and it is further preferred  
 5 that each of Ar and Ar<sup>1</sup> has at least one -COOH and more preferably at least two -COOH groups, for example, dicarboxyphenyl. It is preferred that both Ar and Ar<sup>1</sup> are the same, and more especially that each is 3,5-dicarboxyphenyl.

It is preferred that each of J & J<sup>1</sup> is of Formula (3) and that  
 10 each R<sup>5</sup> is independently selected from H, optionally substituted C<sub>1-4</sub>-alkyl, C<sub>1-4</sub>-alkoxy, Cl, CN, Br, ureido and NHCOR<sup>6</sup>, more preferably H, C<sub>1-4</sub>-alkyl, C<sub>1-4</sub>-alkoxy, ureido and NHCO(C<sub>1-4</sub>-alkyl), and especially H, methyl, methoxy or ureido. When R<sup>5</sup> is not H it is preferably ortho with respect to the azo group (-N=N-).

15 When J and J<sup>1</sup> are of Formula (4) or (5), each T is preferably independently C<sub>1-6</sub>-alkyl, especially C<sub>1-4</sub>-alkyl; the group M may be branched chain alkylene but is preferably straight chain alkylene; W is preferably H, CN, CONR<sup>10</sup>R<sup>11</sup> or COOH; R<sup>6</sup> is preferably H, C<sub>1-4</sub>-alkyl, phenyl or (CH<sub>2</sub>)<sub>1-4</sub>-phenyl, each of which may be unsubstituted or  
 20 substituted, and more especially C<sub>1-4</sub>-alkyl and B is preferably H.

It is preferred that n is 0 so that the two amino groups, NR<sup>1</sup> and NR<sup>4</sup>, are linked by a single group, X, and it is further preferred that X is of Formula (6).

Z is preferably NR<sup>8</sup>R<sup>9</sup>, especially NHC<sub>2</sub>H<sub>4</sub>OH, N(C<sub>2</sub>H<sub>4</sub>OH)<sub>2</sub>, morpholino,  
 25 NH(C<sub>1-6</sub>-alkyl), NH-(CH<sub>2</sub>)<sub>2</sub>-CO<sub>2</sub>H, NHCH<sub>2</sub>C<sub>6</sub>H<sub>4</sub>CO<sub>2</sub>H, mono- or di-carboxyanilino, NHC<sub>6</sub>H<sub>4</sub>SO<sub>3</sub>H or NHCH<sub>2</sub>SO<sub>3</sub>H.

R<sup>1</sup>, R<sup>2</sup>, R<sup>3</sup> and R<sup>4</sup> are preferably each independently H, C<sub>1-4</sub>-alkyl or C<sub>1-4</sub>-hydroxyalkyl, and more especially H.

R<sup>7</sup>, R<sup>8</sup> and R<sup>9</sup> are preferably independently selected from H,  
 30 optionally substituted C<sub>1-6</sub>-alkyl, C<sub>3-4</sub>-alkenyl, optionally substituted phenyl and optionally substituted (CH<sub>2</sub>)<sub>1-4</sub>-phenyl, more preferably from H, allyl, C<sub>1-4</sub>-alkyl, benzyl, C<sub>1-4</sub>-hydroxyalkyl and especially from H, methyl, ethyl and 2-hydroxyethyl. The optional substituents on R<sup>7</sup>, R<sup>8</sup> and R<sup>9</sup> are preferably independently selected from -OH, -SO<sub>3</sub>H and -COOH,  
 35 especially -OH. When R<sup>8</sup> and R<sup>9</sup> together with the nitrogen atom to which they are attached form a 5 or 6 membered ring this is preferably a morpholine or piperidine ring.




The yellow dye is described in greater detail in EP 468,647 the contents of which are incorporated herein by reference.

An especially preferred yellow dye, hereinafter called Dye Y1, is a dye of Formula (2) in which Ar and Ar<sup>1</sup> are both 3,5-dicarboxy-phenyl; J and J<sup>1</sup> are both phen-1,4-ylene; R<sup>1</sup> and R<sup>4</sup> are both H; n is 0; X is of Formula (6) and Z is morpholino, in the ammonium (<sup>+</sup>NH<sub>4</sub>) salt form.

#### Cyan Dye

The cyan dye, in the free acid form, has the structure shown in Formula (9) or the structure shown in Formula (10).

10 (a) Cyan dye of Formula (9)



Formula (9)

wherein:

20 Pc is a metal-free or metal-containing phthalocyanine radical having a valency from 3 to 4;

R<sup>1</sup> is H, optionally alkyl, alkenyl or aralkyl;

R<sup>2</sup> is H, alkyl, alkoxy, halo or optionally substituted amino; and

25 (t+q) is from 3 to 4 inclusive.

In the dye of Formula (9), the COOH group is preferably at the 2-, 3-, 5- or 6-position, especially the 3-position, on the phenyl group.

The metal present in the metal-containing phthalocyanine radical may be any of the metals known to complex with phthalocyanine nuclei, but is preferably scandium, titanium, vanadium, chromium, manganese, iron, cobalt or zinc, more preferably nickel and especially copper. When the metal is tri- or tetra-valent, it may coordinate with chlorine or oxygen, e.g. VO, AlCl<sub>3</sub>, PbCl<sub>2</sub>.

The group R<sup>1</sup> is preferably H; C<sub>3-4</sub>-alkenyl, such as allyl; optionally substituted C<sub>7-11</sub>-aralkyl, especially benzyl; optionally substituted C<sub>1-4</sub>-alkyl, especially methyl, ethyl or hydroxyethyl.

The group R<sup>2</sup> is preferably H; C<sub>1-4</sub>-alkyl, especially methyl; C<sub>1-4</sub>-alkoxy, especially methoxy; chloro; or an amino group, -NX<sup>1</sup>X<sup>2</sup>,

wherein  $X^1$  and  $X^2$  are each independently H, or optionally substituted alkyl or aryl. When  $X^1$  or  $X^2$  is alkyl it is preferably  $C_{1-18}$ -alkyl, more preferably  $C_{1-6}$ -alkyl. When  $X^1$  or  $X^2$  is optionally substituted aryl it is preferably optionally substituted phenyl, especially phenyl or phenyl  
 5 having 1 or 2 substituents selected from  $C_{1-12}$ -alkyl,  $C_{1-12}$ -alkoxy, halo and carboxy. Examples of amino groups,  $NX^1X^2$ , are n-hexylamino, N,N-di-n-butylamino, phenylamino, methylamino, dimethylamino, diethylamino, dipropylamino, ethylamino, n-propylamino, n-butylamino, n-octylamino and 3-carboxyphenylamino.

10 It is especially preferred that  $R^1$  and  $R^2$  are both H.

The value of  $t$  is preferably at least 0.5, more preferably from 0.5 to 2, and especially about 1; the value of  $q$  is preferably greater than or equal to  $t$ , and especially about 3; and  $(t+q)$  is preferably 4. Where the value, of  $(t+q)$  is not integral, the dye of Formula (9) will  
 15 actually comprise a mixture of dyes with different values of  $(t+q)$ .

The cyan dye of Formula (9) is described in greater detail in EP 559309A the contents of which are incorporated herein by reference.


An especially preferred cyan dye, hereinafter called Dye C1, is of Formula (9) wherein Pc is copper phthalocyanine;  $R^1$  and  $R^2$  are both H,  
 20  $t$  is 1 and  $q$  is 3, and the COOH group is in the 3-position, in the ammonium ( $^+NH_4$ ) salt form.

(b) Cyan dye of Formula (10)




25 wherein:


- Pc is a metal-containing phthalocyanine nucleus;  
 $R^1$ ,  $R^2$  and  $R^3$  are each independently H, alkyl, substituted alkyl, alkenyl, substituted alkenyl, aralkyl or substituted aralkyl;  
 30 L is a divalent organic linking group;  
 G is a colourless organic radical substituted by one or two groups selected from COSH and COOH;  
 $(t+q)$  is from 3 to 4 inclusive; and  
 each X independently is carbonyl or a group of formula (11), (12)  
 35 or (13):



(11)



(12)



(13)

wherein:

- 10 each Z independently is  $\text{NR}^4\text{R}^5$ ,  $\text{SR}^6$  or  $\text{OR}^6$ ;  
 each Y independently is H, Cl, Z,  $\text{SR}^7$  or  $\text{OR}^7$ ;  
 each E independently is Cl or CN;  
 $\text{R}^4$ ,  $\text{R}^5$ ,  $\text{R}^6$  and  $\text{R}^7$  are each independently H, alkyl, substituted  
 alkyl, aryl, substituted aryl, aralkyl or substituted aralkyl  
 15 or  $\text{R}^4$  and  $\text{R}^5$  together with the nitrogen atom to which they  
 are attached form a 5 or 6 membered ring.

The cyan dye of Formula (10) is preferably free from COSH groups  
 and also preferably contains at least as many  $-\text{COOH}$  groups as  $-\text{SO}_3\text{H}$   
 groups.

- 20 The metal present in the metal-containing phthalocyanine nucleus  
 may be any of the metals known to be found in complex association with  
 phthalocyanine nuclei, but is preferably selected from Sc, Ti, V, Cr,  
 Mn, Fe, Co and Zn, and more especially from Ni and Cu. When the metal  
 is tri- or tetra-valent, it may carry one or two chlorine atoms.

- 25  $\text{R}^1$ ,  $\text{R}^3$ ,  $\text{R}^6$  and  $\text{R}^7$  are preferably each independently selected from  
 H,  $\text{C}_{1-4}$ -alkyl and hydroxy- $\text{C}_{1-4}$ -alkyl, more preferably H, and  $\text{C}_{1-4}$ -alkyl.  
 It is particularly preferred that  $\text{R}^1$  and  $\text{R}^3$  are H.

The group  $\text{R}^2$  is preferably H,  $\text{C}_{1-4}$ -alkyl, such as methyl or  
 hydroxy- $\text{C}_{1-4}$ -alkyl, such as hydroxyethyl.

- 30 The groups  $\text{R}^4$  and  $\text{R}^5$  are preferably each independently H, or  
 optionally substituted  $\text{C}_{1-4}$ -alkyl, preferred substituents being  $-\text{COOH}$ ,  $-\text{SO}_3\text{H}$  or, more especially,  $-\text{OH}$ . Where  $\text{R}^4$  and  $\text{R}^5$  together with the N atom  
 to which they are attached form a 5- or 6-membered ring, this is  
 preferably morpholinyl or piperidinyl.

- 35 When X is of Formula (12), Z is preferably attached to the C atom  
 between the two ring N atoms and Y is preferably para with respect to Z.  
 However, X is preferably of Formula (11).

Z is preferably  $\text{NR}^4\text{R}^5$ , especially  $\text{C}_{1-4}$ -hydroxyalkylamino such as hydroxyethylamino, or OH.

The value of q is preferably greater than or equal to the value of t and (t+q) is preferably 4.

5 The identity of the divalent organic linking group L is not critical but is preferably  $\text{C}_{2-6}$ -alkylene, such as eth-1,2-ylene, or optionally substituted phenylene.

G is preferably substituted alkyl, especially  $\text{C}_{1-4}$ -substituted alkyl; substituted aralkyl, especially substituted phenyl- $(\text{C}_{1-3}$ -alkyl);  
 10 substituted phenyl; or substituted naphthyl wherein at least one of the substituents is selected from  $-\text{COOH}$ ,  $-(\text{CH}_2)_n-\text{COOH}$ ,  $-\text{O}-(\text{CH}_2)_n-\text{COOH}$ ,  $-\text{S}-(\text{CH}_2)_n-\text{COOH}$  and  $-\text{NR}^8(\text{CH}_2)_n-\text{COOH}$  in which n is an integer from 1 to 6 and  $\text{R}^8$  is H or  $\text{C}_{1-4}$ -alkyl. The group G is preferably carboxyphenyl or dicarboxyphenyl.

15 The cyan dye of Formula (10) is described in greater detail in EP 468,649 the contents of which are incorporated herein by reference.

The present invention relates not only to an ink containing a colorant comprising a black and a yellow and/or a cyan dye as hereinbefore defined in the free acid form, but also in the form of  
 20 salts, preferably alkali metal, ammonium or substituted ammonium salts, and especially salts derived from ammonia or a volatile amine. Furthermore, although any amino groups in the formulae for the dyes are shown in neutral form, these groups may exist as substituted ammonium cations, especially if the dye is capable of existing in zwitterionic  
 25 form.

In the salt form, especially as alkali metal, ammonium or substituted ammonium salts or mixtures thereof, at pH 7.5 and above, especially from pH 7.5 to 9.0, the dyes have good solubility in water, often up to 10% or more, by weight. However, the water-solubility drops  
 30 sharply below pH 7.5 and below pH 6 they are generally insoluble in water and aqueous media.

The cation in the ammonium or substituted ammonium salts is preferably a quaternary ammonium group of the formula  $^+\text{NH}_p\text{Q}_q$  in which p and q are each independently integers from 0 to 4 provided that  
 35  $(p+q) = 4$ , and each Q independently is an organic radical; or 2 or 3 Qs, together with the N atom to which they are attached, form a heterocyclic ring and any remaining Qs are  $\text{C}_{1-4}$ -alkyl.

Organic radicals represented by Q are preferably C<sub>1-4</sub>-alkyl, especially methyl; aryl, especially phenyl; or aralkyl, especially benzyl. Heterocyclic rings formed by NQ<sub>4</sub> are preferably 5- or 6-membered heterocycles, such as morpholinyl, piperidinyl or pyridyl.

5        Examples of quaternary ammonium groups of formula <sup>+</sup>NH<sub>p</sub>Q<sub>q</sub> are <sup>+</sup>NH<sub>3</sub>(CH<sub>3</sub>), <sup>+</sup>NH<sub>2</sub>(CH<sub>3</sub>)<sub>2</sub>, <sup>+</sup>NH(CH<sub>3</sub>)<sub>3</sub>, <sup>+</sup>N(CH<sub>3</sub>)<sub>4</sub>, <sup>+</sup>N(CH<sub>2</sub>CH<sub>3</sub>)<sub>4</sub>, N-methyl pyridinium, N,N-dimethyl piperidinium and N,N-dimethylmorpholinium.

10        The dyes can be converted into their ammonium or substituted ammonium salts by dissolving in water, as a salt with an alkali metal, acidifying the solution with a mineral acid, adjusting the pH to 9-9.5 with ammonia or an amine or a quaternary ammonium hydroxide and removing alkali metal salt by dialysis.

15        The inks may be prepared by dissolving the dyes, individually or mixed together to form the colorant, in water alone and optionally adding a water-soluble organic liquid or in an ink medium comprising water and a water-soluble organic liquid. Other ingredients can be added to the ink either before or after the dyes. The ink is preferably filtered prior to use in ink jet printing in order to remove any undissolved matter which can cause blockages in the nozzles of an ink  
20        jet printer.

The invention is further illustrated by the following Examples in which all parts and percentages are by weight unless otherwise indicated.

#### Examples

25        Appropriate amounts of the dyes identified in Table 1 were dissolved in the identified ink media in order to prepare the Inks 1 to 11 identified in Table 1 each ink containing 2.5% colorant. The inks were then assessed for printing quality by printing onto an acid paper (Ac, Xerox 4024 DP) and an alkaline paper (Alk, Hammermill Fore DP3V-2)  
30        using a Hewlett-Packard Deskjet printer. The parameters measured were print density (PD), luminance (L), colour space coordinates (A & B), wet-fastness (WF (%)), light-fastness (LF (Blue scale rating)) and visual appearance (VA), the latter to detect the presence or absence and extent of bronzing. The components of the ink media (Med) were water  
35        (W), 2-pyrrolidone (P) and sulpholane (S) and the proportions (prop) were by weight.

Table 1

| <u>Ink</u> | <u>Dye</u> | <u>%</u> | <u>Med &amp; Prop</u> | <u>Paper</u> | <u>PD</u> | <u>L</u> | <u>A</u> | <u>B</u> | <u>WF</u> | <u>LF</u> |
|------------|------------|----------|-----------------------|--------------|-----------|----------|----------|----------|-----------|-----------|
| 5          | 2          | B1 95    | W:P 90:10             | Ac           | 25.5      | 27.1     | 2.4      | -5.6     | 97        | 3         |
|            |            | C1 5     |                       | Alk          | 14.5      | 34.5     | 3.8      | -7.7     | 94        | 3         |
|            | 3          | B1 90    | W:P 90:10             | Ac           | 23.2      | 28.4     | 2.8      | -4.9     | 97        | 3         |
|            |            | C1 10    |                       | Alk          | 16.5      | 32.7     | 3.2      | -7.8     | 94        | 3         |
| 10         | 4          | B1 95    | W:P 90:10             | Ac           | 26.5      | 26.8     | 1.7      | -3.8     | 98        | 3         |
|            |            | Y1 5     |                       | Alk          | 14.7      | 34.5     | 2.9      | -6.1     | 92        | 3         |
|            | 5          | B1 90    | W:P 90:10             | Ac           | 27.5      | 26.5     | 1.1      | -3.5     | 98        | 3         |
|            |            | Y1 10    |                       | Alk          | 17.8      | 32.2     | 2.2      | -4.4     | 91        | 3         |
|            | 6          | B1 95    | W:S 95:5              | Ac           | 28.7      | 25.6     | 3.5      | -4.5     | 96        | 3         |
|            |            | C1 5     |                       | Alk          | 18.2      | 31.3     | 4.2      | -7.3     | 90        | 3         |
| 15         | 7          | B1 90    | W:S 95:5              | Ac           | 28.9      | 25.5     | 3.2      | -4.7     | 96        | 3         |
|            |            | C1 10    |                       | Alk          | 16.8      | 32.3     | 3.9      | -8.3     | 91        | 3         |
|            | 8          | B1 95    | W:S 95:5              | Ac           | 27.9      | 26.1     | 2.7      | -4.0     | 99        | 3         |
|            |            | Y1 5     |                       | Alk          | 22.0      | 29.1     | 4.0      | -4.8     | 84        | 3         |
| 20         | 9          | B1 90    | W:S 95:5              | Ac           | 28.6      | 26.0     | 2.5      | -3.0     | 98        | 3         |
|            |            | Y1 10    |                       | Alk          | 20.0      | 30.5     | 3.1      | -4.5     | 84        | 3         |
|            | 10         | B1 90    | W:P 90:10             | Ac           | 27.3      | 26.4     | 1.5      | -4.3     | 97.5      | 3         |
|            |            | C1 5     |                       |              |           |          |          |          |           |           |
| 25         | 11         | B1 85    | W:P 90:10             | Ac           | 26.7      | 27.0     | 0.6      | -2.8     | 97.5      | 3         |
|            |            | Y1 15    |                       |              |           |          |          |          |           |           |

Notes:

Med &amp; Prop is ink medium and proportions by weight

W is water

30 S is sulpholane

Tr Br is trace bronzing

No Br is no bronzing

Claims

1. An ink comprising a solution of a colorant in an aqueous medium, the colorant comprising at least 80% by weight of a water-soluble black dye of Formula (1) as herebefore defined and up to 20% of one or both of two other water-soluble dyes, one of which is a yellow dye of the Formula (2) as hereinbefore defined and the other is a cyan dye of Formula (9) as hereinbefore defined.
2. An ink according to Claim 1 wherein the colorant comprises from 90% to 98% of the black dye and from 10% to 2% of the yellow dye.
3. An ink according to Claim 1 wherein the colorant comprises from 90% to 95% of the black dye and from 10% to 5% of the cyan dye.
4. An ink according to any one of Claims 1 to 3 wherein the black dye is the dye of Formula (1) wherein W is 3-COOH, X is 5-COOH and Y, Z and R are H, in the ammonium salt form.
5. An ink according to Claim 1 or Claim 2 wherein the yellow dye of Formula (2) in which Ar and Ar<sup>1</sup> are both 3,5-dicarboxyphenyl; J and J<sup>1</sup> are both phen-1,4-ylene; R<sup>1</sup> and R<sup>4</sup> are both H; and n is 0, in the ammonium salt form.
6. An ink according to Claim 1 or Claim 3 wherein the cyan dye is of Formula (9) wherein Pc is copper phthalocyanine; R<sup>1</sup> and R<sup>2</sup> are both H, t is 1 and q is 3, and the COOH group is in the 3-position, in the ammonium salt form.

**Patents Act 1977**  
**Examiner's report to the Comptroller under Section 17**  
**(The Search report)**

15.

Application number  
 GB 9402090.6

**Relevant Technical Fields**

- (i) UK Cl (Ed.M) C4P (PM)  
 (ii) Int Cl (Ed.5) CO9B 67/22; CO9D 11/00; CO9D 11/02

**Databases (see below)**

- (i) UK Patent Office collections of GB, EP, WO and US patent specifications.

- (ii) ONLINE DATABASES: CA, WPI, CLAIMS

Search Examiner  
 A J RUDGE

Date of completion of Search  
 19 MAY 1994

Documents considered relevant  
 following a search in respect of  
 Claims :-  
 ALL

**Categories of documents**

- X:** Document indicating lack of novelty or of inventive step.      **P:** Document published on or after the declared priority date but before the filing date of the present application.
- Y:** Document indicating lack of inventive step if combined with one or more other documents of the same category.      **E:** Patent document published on or after, but with priority date earlier than, the filing date of the present application.
- A:** Document indicating technological background and/or state of the art.      **&:** Member of the same patent family; corresponding document.

| Category | Identity of document and relevant passages | Relevant to claim(s) |
|----------|--|----------------------|
| A        | EP 0559309 A2 (ZENECA)*                    | ALL                  |
| A        | EP 0565286 A1 (ICI)*                       | ALL                  |
| A        | EP 0534634 A1 (HEWLETT-PACKARD)*           | ALL                  |
| A        | EP 0494523 A1 (ICI)*                       | ALL                  |
| A        | EP 0468649 A1 (ICI)*                       | ALL                  |
| A        | EP 0468648 A1 (ICI)*                       | ALL                  |
| A        | EP 0468647 A1 (ICI)*                       | ALL                  |
| A        | EP 0356080 A2 (ICI)*                       | ALL                  |
| A        | US 4963189 (HEWLETT-PACKARD)*              | ALL                  |
|          | * see whole document                       |                      |

Databases: The UK Patent Office database comprises classified collections of GB, EP, WO and US patent specifications as outlined periodically in the Official Journal (Patents). The on-line databases considered for search are also listed periodically in the Official Journal (Patents).